




Anicla infecta Green Cutworm Moth



FAMILY: Noctuidae SUBFAMILY: Noctuinae TRIBE: Noctuini

TAXONOMIC COMMENTS: One of fifteen species in this genus that occur in North America north of Mexico (Lafontaine and Schmidt), six of which have been recorded in North Carolina

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Wagner et al. (2011) list fields, waste lots, croplands, marshes, wet meadows and other open habitats. Our records additionally include mesic forests and dry woodlands.

FOOD: Larvae feed on grasses as well as a wide range of forbs, including numerous crop species (Wagner et al., 2011). Pogue (2006) specifically lists beets, tobacco, plantain, purslane, and clover. We do not have any feeding records in North Carolina.

OBSERVATION_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 S5

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: