




Xestia c-nigrum Lesser Black-letter Dart Moth



High counts of:
 1 - Stokes - 1998-08-18
 1 - Ashe - 2007-08-23
 1 - Iredell - 2018-09-07

Status	Rank
NC	US
NC	Global



FAMILY: Noctuidae SUBFAMILY: Noctuinae TRIBE: Noctuini
 TAXONOMIC_COMMENTS:

FIELD GUIDE DESCRIPTIONS: Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT:

FOOD: Larvae are polyphagous, feeding on a wide range of plants, including many cultivated species. Reported hosts include alfalfa, barley, blueberry, chokeberry, clover, corn, cranberry, dandelion, flax, goldenrod, grape, lettuce, maple, nettle, oats, pear, sunflower, tobacco, tomato, viburnum, and wheat (Crumb, 1932; Robinson et al., 2010).

OBSERVATION_METHODS:

NATURAL HERITAGE PROGRAM RANKS:

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: