



Schinia sordidus Sordid Flower Moth



High counts of:

- 2 - Pender - 1995-08-30
- 2 - Columbus - 2009-09-18
- 1 - Brunswick - 1995-08-28

Status Rank
NC US NC Global



FAMILY: Noctuidae SUBFAMILY: Heliothinae TRIBE:

TAXONOMIC COMMENTS: One of 126 species in this genus that occur in North America (Lafontaine and Schmidt, 2010, 2011), the majority of which occur in the West; 25 have been recorded in North Carolina.

FIELD GUIDE DESCRIPTIONS:

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Our records all come from dry, open sand barrens, including both open maritime scrub, xeric Carolina Bay rims, and inland sandhills.

FOOD: Reported hosts include Sandhill Golden-aster (*Pityopsis pinifolia*) and Slender Scratch-daisy (*Croptilon divaricatum*) (Wagner et al., 2011).

OBSERVATION_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G4? S2S3

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: