




Schinia siren Aluring Schinia Moth



High counts of:
 17 - Brunswick - 1994-09-12
 14 - Brunswick - 1994-09-05
 2 - Carteret - 2021-09-27

Status	Rank
NC	US
NC	Global



FAMILY: Noctuidae SUBFAMILY: Heliiothinae TRIBE:
 TAXONOMIC COMMENTS: One of 126 species in this genus that occur in North America (Lafontaine and Schmidt, 2010, 2011), the majority of which occur in the West; 25 have been recorded in North Carolina.

FIELD GUIDE DESCRIPTIONS:
 ONLINE PHOTOS:
 TECHNICAL DESCRIPTION, ADULTS:
 TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT:

FOOD: Reported hosts include camphorweed (<i>Heterotheca spp.</i>) and <i>Verbesina</i> (Heppner, 2007; Beadle & Leckie, 2018). We do not have any larval records in North Carolina.

OBSERVATION_METHODS:

NATURAL HERITAGE PROGRAM RANKS: GNR [S3S4]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: