




# *Schinia florida* Primrose Moth



FAMILY: Noctuidae SUBFAMILY: Heliiothinae TRIBE:

TAXONOMIC COMMENTS: One of 126 species in this genus that occur in North America (Lafontaine and Schmidt, 2010, 2011), the majority of which occur in the West; 25 have been recorded in North Carolina.

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Wagner et al. (2011) list fields, roadsides, waste places, grasslands, prairies, and coastal communities as habitats used by this species. Our records all come from the Mountains, including from wet meadows, mountain bogs, pastures, and high elevation fields.

FOOD: Larvae feed on Biennial Beeblossom (*Oenothera gaura*) and Evening Primroses (*Oenothera* sp.) (Wagner et al., 2011).

OBSERVATION\_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 S2S4

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: