
Schinia lucens Leadplant Flower Moth



FAMILY: Noctuidae SUBFAMILY: Heliothinae TRIBE:

TAXONOMIC_COMMENTS: One of 126 species in this genus that occur in North America (Lafontaine and Schmidt, 2010, 2011), the majority of which occur in the West; 25 have been recorded in North Carolina.

FIELD GUIDE DESCRIPTIONS: Covell (1984)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: The heart of this species' range is in the Tall-grass Prairie Region of the upper Mid-west. Peripheral populations occur in California, Arizona, and along the Atlantic and Gulf Coastal Plains (Moth Photographers Group)

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT:

FOOD: Larvae reportedly feed on species of <i>Amorpha</i>, including False Indigo-bush (<i>A. fruticosa</i>) and Leadplant (<i>A. canescens</i>) (Wagner et al., 2011).

OBSERVATION METHODS:

NATURAL HERITAGE PROGRAM RANKS: G4 SNR [SH]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: