




Stereomita andropogonis No common name



High counts of:
 1 - Onslow - 2019-09-21
 1 - Orange - 2022-09-12
 1 - Orange - 2023-09-04

Status		Rank	
NC	US	NC	Global



FAMILY: Gelechiidae SUBFAMILY: Anomologinae TRIBE:
 TAXONOMIC_COMMENTS:

FIELD GUIDE DESCRIPTIONS:

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS: The following is based on Braun's (1922) original description. The head is whitish straw-colored and smooth. The labial palp is also straw-colored, with a dark brown patch near the apex of the second segment outwardly, and a dark brown annulus around the middle of the third segment. The antenna is slightly shorter than the length of the forewing and is pale ochereous, with a narrow brown annulus at the base of each segment. There are four broad blackish bands on the apical half of the stalk that are separated from one another by two or three pale segments. The forewing is pale ochereous and dusted with dark brown scales. These are most dense on the costal and dorsal margins, and in some individuals may resemble diffuse longitudinal streaks. A series of indistinct brownish dots are present along the termen, and a transverse brownish spot is often present in the cilia opposite the extreme apex. The cilia are brownish overall, except for more ochereous coloration along the costa before the apex. The hindwing is pale brown and the cilia is ochereous, with a faint reddish tinge. The legs are ochereous and dusted with brown. The posterior tibia has rough hairs above, and in the middle below. Individuals that are heavily dusted with dark brown scales resemble *Batrachedra busiris*. The black-banding on the antenna may be the easiest way to distinguish these. On *S. andropogonis* the bars begin near the middle of the antenna, while on *B. busiris* they are confined to the apical fourth. The palp also has a conspicuous dark ring on the third segment of *S. andropogonis* that is absent on *B. busiris*.

DISTRIBUTION: *Stereomita andropogonis* is found in eastern North America, where the range extends from Massachusetts, New York, Quebec, and Ontario, then southward to the Gulf Coast and westward to central Texas, Oklahoma, and Illinois. As of 2021, we have a single record from near the coast.

FLIGHT COMMENT: The adults have been observed from June through September in areas outside of North Carolina. As of 2021, our only record is from September.

HABITAT: Little Bluestem is the only known host. This species is widespread in the state in a variety of open, sunny habitats or open woodlands. It generally prefers drier habitats where rank, weedy vegetation is suppressed. *Stereomita andropogonis* is associated with prairie habitats in many areas of its range (e.g., Metzler, 1997), which likely reflects that fact that Little Bluestem is an important member of many prairie communities.

FOOD: The only known host is Little Bluestem (*Schizachyrium scoparium*).

OBSERVATION_METHODS: The adults are attracted to lights. Braun (1922) observed the moths flying around the tops of stems and flower buds in early morning and in the evening. They characteristically landed with the head pointing downward. They were inactive during the middle of the day and rested among the basal leaves. More information is needed on host plant use and the larval ecology and life history of this species.

NATURAL HERITAGE PROGRAM RANKS: GNR S1S2

STATE PROTECTION:

COMMENTS: We currently do not have sufficient information on the distribution and abundance of this species to assess its conservation status within the state.