




*Chionodes pereyra* No common name



FAMILY: Gelechiidae SUBFAMILY: Gelechiinae TRIBE: Gelechiini

TAXONOMIC COMMENTS: The genus *Chionodes* is the most species rich genus of gelechiid moths in the Western Hemisphere, with 187 recognized species. Our knowledge of the diverse array of species in North America is largely due to the monumental work of Hodges (1999), who spend decades working on the group and described 115 new species (Powell and Opler, 2009). Many exhibit substantial variation within species and have drab coloration, typically with brown, dark gray, or blackish patterning on the forewings. These can only be confidently identified by examining secondary sexual characteristics and/or the genitalia of one or both sexes. Others are more boldly marked and can be identified by wing patterning. Many of our state records are based on Hodges (1999) database of over 19,000 specimens that he examined from major collections in the US. These include North Carolina specimens that he collected mostly from Highlands, and from a few other areas within the state.

FIELD GUIDE DESCRIPTIONS: The larvae are known to feed on oaks, but very little is known about the larval life history.

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS: Hodges (1999)

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS: The labial palp is pale luteous and suffused with light brown in the brush. The second segment is irrorate with blackish fuscous exteriorly, while the third segment is blackish fuscous. The head is ochreous-white to sordid white, with the scales brown to fuscous-tipped. The antenna is blackish fuscous with light-brown annulations. The ground color of the thorax and forewing is light-brown. A moderately large, blackish-fuscous spot is on the costa at the basal third. Elongated blackish-fuscous spots are present on the fold just before the middle of the wing, in the center of the cell, and at the end of cell. The apical third of the wing is fuscous and bisected by a pale, outwardly angulate, transverse fascia at the apical fourth. The cilia are light yellowish fuscous, mixed with fuscous. The hindwing is pale gray basally, and shades to dark fuscous apically. The legs are ochreous-white and suffused and banded with fuscous. The abdomen is light fuscous above, with the posterior edges of the segments narrowly bordered with ochreous-white to sordid white. This species is best identified using genitalia.

DISTRIBUTION: *Chionodes pereyra* is found from Massachusetts and vicinity southward to southern Florida, and westward to eastern Texas, Oklahoma, Kansas, Illinois, and Michigan. As of 2021, almost of our records are from Highlands in Macon County, where Hodges (1999) collected extensively. We have a single records from the coast.

FLIGHT COMMENT: Hodges (1999) reported the flight season to extend from mid-March to mid-September, with most records from June and July. As of 2021, we have records from January through December, with a seasonal peak in July.

HABITAT: Populations are associated with hardwood forests with oaks.

FOOD: The larvae feed on members of the red oak group (Hodges, 1999). The known hosts include Scarlet Oak (*Quercus coccinea*) and Northern Red Oak (*Q. rubra*).

OBSERVATION\_METHODS: The adults come to lights, particularly on rainy nights (Hodges, 1999).

NATURAL HERITAGE PROGRAM RANKS: GNR SU

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: We currently do not have sufficient information on the distribution and abundance of this species to assess its conservation status.