



*Glyphipterix saurodonta* No common name



Status		Rank	
NC	US	NC	Global



FAMILY: Glyphipterigidae SUBFAMILY: TRIBE:  
 TAXONOMIC\_COMMENTS:

FIELD GUIDE DESCRIPTIONS:

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS: Heppner (1985).

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS: The following description is based on the descriptions by Meyrick (1913) and Heppner (1985). The head and thorax are bronzy-fuscous, with a white line above the eye. The palp has three alternating bands of fuscous and white, and the apical portion of the terminal joint is fuscous with white lateral lines. The antenna is fuscous dorsally. The forewing ground color is dark bronzy-gray to fuscous and there is a very narrow white line along the anal margin. The costa has six silvery-white streaks that are edged with dark fuscous. The first two are slightly oblique and reach nearly half way across the wing, with the first at one-third the length from the wing base to the apex, and the second at the middle. The remaining four are shorter, with a pair immediately behind the second streak, and another pair just before the apex. The inner margin has a posteriorly oblique silvery-white streak that begins at around three-fourths and extends to where it almost reaching the apex of the second costal streak. This is followed by a shorter, silvery-white, dark-edged streak just before the tornus that runs more perpendicular to the inner margin. The space between this and the preceding dorsal streak is whitish-tinged. There are three silvery-metallic dots on the lower half of the termen (last two sometimes confluent), and two others obliquely placed in the disc above these. There is also a silvery-metallic dot on the termen beneath the apex, which is suffused with blackish coloration. The fringe is gray, with the basal half rather dark fuscous. It is indented with white on the subapical dot and has a blackish hook at the apex. The hindwing is dark gray with gray cilia.

<i>Glyphipterix chambersi</i> is similar but has the white line above the eye reduced and normally has five (rarely six) white costal streaks rather than six as seen in <i>G. saurodonta</i>. It is also found at more northern latitudes (northern Kentucky and southern Canada), with flight records all from August. Heppner (1985) noted that when a sixth streak is present on <i>G. chambersi</i> that one is minute and lacks the fuscous border. Rob Van Epps discovered a specimen from North Carolina that we have provisionally assigned to <i>G. saurodonta</i> rather than <i>G. chambersi</i> based on its well-developed eye stripe, the presence of six well-developed white costal streaks, the date of collection (late October) and the geographic location.

DISTRIBUTION: <i>Glyphipterix saurodonta</i> is rare in collections and is only known from Toronto, Vermont, New York, West Virginia, and eastern Tennessee (Heppner, 1985; Moth Photographers Group). As of 2023, we have a single record from Mecklenburg County.

FLIGHT COMMENT: The known flight records extend from mid-August through October (Heppner, 1985; iNaturalist). As of 2023, our one record is from late-October.

HABITAT: The preferred habitats are poorly documented. Our one specimen as of 2023 was collected by sweeping in a field near some deciduous trees.

FOOD: The host plants are unknown.

OBSERVATION\_METHODS:

NATURAL HERITAGE PROGRAM RANKS: GNR [S1S2]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: This species appears to be rare throughout its range. As of 2023, we have a single record for the state.