



*Epiblema mandana* No common name



High counts of:  
 9 - Durham - 2022-08-02  
 2 - Mecklenburg - 2023-07-28  
 1 - Buncombe - 2021-06-28

Status		Rank	
NC	US	NC	Global



FAMILY: Tortricidae SUBFAMILY: TRIBE:  
 TAXONOMIC\_COMMENTS:

FIELD GUIDE DESCRIPTIONS:

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS: Wright and Gilligan (2023)

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS: The following description is based on that of Wright and Gilligan (2023). In this species the vertex and thorax are brown, while the frons is tan. The antenna is brown with conspicuous black marks on the pedicel and the scape. The third segment of the labial palp is black, and the lateral surface of the second segment is pale brown with a diffuse black mark near the dorsal margin. The lateral surface of first segment has a faint black dot. The forewing is medium brown and often has grayish tints. The fasciate markings seen on many other *Epiblema* species are absent to barely discernable, and when present are represented as slightly darker shades of brown. The dorsal (interfascial) spot is slightly paler than the adjacent sub-basal and median areas and sometimes has faint orange-brown tints, but is never sharply defined. The ocellus is whitish and conspicuous, and the costal strigulae are white and separated by orange-brown lines that extend from the costa towards the termen. Strigula 7 has an associated prominent gray stria that extends to the termen, which has a narrow salt-and-pepper colored band that extends from the tornus to the apex. The hindwing is uniformly gray brown.

This species is mostly easily confused with *E. abruptana*, but the latter typically has black fasciate markings on the forewing versus a rather uniformly medium-brown coloration for *E. mandana*. In addition, the black terminal tip of the palp is usually easily seen on *E. mandana* when viewed from above, versus being hidden or barely visible on *E. abruptana*. Both species have a black mark at the base of the antenna, but it is more pronounced on *E. mandana*. Some specimens may need to be dissected to examine genitalia.

DISTRIBUTION: Wright and Gilligan (2023) identified specimens from Alabama, Mississippi, Arkansas, Kansas, Kentucky, Louisiana, Maryland, Ohio, South Carolina, Tennessee, Texas and West Virginia. As of 2024, our records are all from the Piedmont and lower-elevations in the Blue Ridge.

FLIGHT COMMENT: Specimens that were identified by Wright and Gilligan (2023) from different areas of the range were collected from April through August. As of 2024, our records extend from late-May through early-August.

HABITAT: Our records are mostly from fragmented landscapes that have mixtures of forests, pastures, fields, residential developments and roadway corridors.

FOOD: The larval host plants are undocumented.

OBSERVATION\_METHODS: The adults are attracted to lights.

NATURAL HERITAGE PROGRAM RANKS: GNR[S3S4]

STATE PROTECTION:

COMMENTS: This species may be more common than our records suggest due to confusion with sibling species. More information is needed on its distribution and host use before we can accurately assess its conservation status.