



Dichrorampha leopardana None



FAMILY: Tortricidae SUBFAMILY: Olethreutinae TRIBE: Grapholitini
TAXONOMIC_COMMENTS:

FIELD GUIDE DESCRIPTIONS:

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS: Busck (1906a)

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS: This is a distinctively patterned *Dichrorampha* with bold leopard-like patterning on the forewings, which is the basis for the specific epithet. The following is based in part on the description by Busck (1906a). The head and thorax are bright orange to light yellowish brown with blackish barring, while the labial palp is light yellow, with the tip of the brush and apical joint black. The forewing ground is also bright orange to light yellowish brown and is overlain with a mixture of brownish-black to blackish blotches and streaks. These are best developed on the dorsal two-thirds of the wing and form a reticulated pattern. The costa has a series of dark bands on the basal half that are replaced by shorter streaks on the apical half, along with variably expressed whitish strigulae along the margin. The apical third has two leadened-gray striae that extend from the costa obliquely towards the termen. A weakly formed ocellus is present near the middle of the wing at around three-fourths that has two or three horizontal black dashes, and the termen has four black spots near the middle. The fringe on both the front and hindwing is brown with a pale line near the base, and the hindwing is dark brown.

DISTRIBUTION: *Dichrorampha leopardana* is restricted to eastern North America and has been found from New Jersey and Pennsylvania westward through southern Ontario and the Great Lakes region to Illinois, Wisconsin, Iowa, and eastern Nebraska. The range extends southward to southern Indiana, Ohio, Kentucky, northern Virginia and western North Carolina. As of 2022, we have records from three counties in the Blue Ridge that may represent a southern isolate that is disjunct from the main group farther north.

FLIGHT COMMENT: The adults fly from May through September in different areas of the range. As of 2022, we have records that extend from mid-May to early September.

HABITAT: Local populations are often found in deciduous forests, forest edges, and fields that support the host plants. We have records from mesic forests and a high-elevation grassy bald.

FOOD: The known larval hosts include White Snakeroot (*Ageratina altissima*) and a crownbeard (*Verbesina* sp.; Godfrey et al., 1987; Heinrich, 1926).

OBSERVATION_METHODS: The adults are attracted to lights and the leaf folds can be found on the host plants.

NATURAL HERITAGE PROGRAM RANKS: GNR [S2-S3]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: This species appears to reach the southern limit of its range in western North Carolina and is generally uncommon within the state.