


Argyrotaenia kimballi Kimball's Leafroller Moth



FAMILY: Tortricidae SUBFAMILY: Tortricinae TRIBE: Archipini

TAXONOMIC_COMMENTS: The genus *Argyrotaenia* contains approximately 100 described species, with most occurring in Nearctic and Neotropical regions. Thirty-six species are currently recognized in North America.

FIELD GUIDE DESCRIPTIONS:

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS: According to Obratzsov (1961) *kimballi* differs from *velutinana* in having the basal area of forewings much paler along the costal margin and in the costal spot between the median fascia and wing apex consisting of three more or less distinctly separated parts. In both sexes, *kimballi* has the hind wings much paler than does *velutinana*.

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT:

FOOD: The larval hosts are undocumented (Brown et al., 2008).

OBSERVATION_METHODS:

NATURAL HERITAGE PROGRAM RANKS:

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: