



*Syllonoma longipalpana* No common name



| Status |    | Rank |        |
|--------|----|------|--------|
| NC     | US | NC   | Global |



FAMILY: Tortricidae SUBFAMILY: Tortricinae TRIBE: Sparganothini  
TAXONOMIC\_COMMENTS:

FIELD GUIDE DESCRIPTIONS:

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: The three known localities are situated along 140 km of coastal wetland and beach habitat (Powell and Brown, 2012). We have at least one specimen from an area with extensive Longleaf Pine savannas but no marshes or tidewater scrub.

FOOD: The larval food plants are unknown but members of the Asteraceae seem likely (Powell and Brown, 2012)

OBSERVATION\_METHODS:

NATURAL HERITAGE PROGRAM RANKS: [GNR] S1S2

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: This is rarely collected species that Powell and Brown (2012) knew from only three sites along the coasts of North and South Carolina.