
## Adoneta pygmaea No common name



FAMILY: Limacodidae SUBFAMILY: TRIBE:

TAXONOMIC\_COMMENTS: The genus <i>Adoneta</i> contains four recognized species in North America, with all but one occurring in North Carolina.

FIELD GUIDE DESCRIPTIONS:

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS: Grote and Robinson (1868)

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS: <i>Adoneta pygmaea</i> is the smallest of our slug moths, with a wing expanse of only 14-15 mm (Grote and Robinson, 1868). The head, thorax, palps, and antennae vary from reddish or orangish-tan to a darker reddish-brown. The thorax is covered with long hairs, and the legs tend to be a slightly darker shade, with the front and insides more blackish. The forewing is rounded -- with a ground color typically concolorous with the head and thorax &ndash; and is marked with two whitish lines. The first extends from the inner margin at about one-third the length from the base to the tornus. It curved slightly posteriorly before terminating near the middle of the wing. The second extends from the costa at about four-fifths and parallels the termen before ending near the middle of the wing. The fringe tends to be concolorous with the wing in light forms and dark brown in the darker forms. The hindwing is similar in color to the forewing, but of a lighter shade.

DISTRIBUTION: <i>Adoneta pygmaea</i> is primarily found in Texas, with additional records known from southern Oklahoma and western Louisiana. An isolated record is also known for southeastern Georgia. As of 2023, we have two records from the Sandhills.

FLIGHT COMMENT: The adults have been observed from April through November, with a seasonal peak typically in August and September. As of 2023, we have records from late July and early August.

HABITAT: Our records are from the Sandhills, one near the interface of a xeric Longleaf Pine forest and a headwater riparian hardwood forest and the other from an old farm located on a flat.

FOOD: The larval host plants are undocumented.

OBSERVATION METHODS: The adults are attracted to lights.

NATURAL HERITAGE PROGRAM RANKS: GNR [S1]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: This species appears to be rare in North Carolina. We have only two specimens from the Sandhills, with the nearest known record from southeastern Georgia. We are uncertain if a viable population exists in the Sandhills.