




Dicymolomia julianalis Julia's Dicymolomia



High counts of:
 5 - Durham - 2021-03-27
 5 - Bladen - 2021-03-31
 5 - Madison - 2021-02-25

Status		Rank	
NC	US	NC	Global



FAMILY: Crambidae SUBFAMILY: Glaphyriinae TRIBE: [Glaphyriini]
 TAXONOMIC_COMMENTS:

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012)
 ONLINE PHOTOS:
 TECHNICAL DESCRIPTION, ADULTS: Munroe (1972)
 TECHNICAL DESCRIPTION, IMMATURE STAGES: Munroe (1972)

ID COMMENTS: This is a distinctively marked and shaped moth with large eyes and rather complex patterning. Note the large whitish area in the median area that is peppered with darker scales, the rusty brown region on the apical two-thirds that shades to a lighter yellowish brown basally, and the single row of black dots on the outer margin of the hindwing. The following description is based in part on the descriptions by Forbes (1923) and Monroe (1972). The labial palp is upturned and has more or less rough hairs. The head is whitish above, and the antenna is light brown with obscure darker annulations. The thorax and base of the wings are more or less concolorous and interrupted by a white streak that extends backwards along the side of the thorax. One or two shorter and less conspicuous white streaks are usually evident between this streak and the costal margin. The forewing has a rusty brown region on the apical third and just before one-half the wing length. The two are interrupted by a broad zone of white that is heavily dusted with dark brown. The rusty brown region just before one-half shades into bright yellow to light yellowish brown towards the base. A region of black dusting is usually present near the apex on the costal third, and a whitish terminal line is usually evident at the wing tip. The hindwing is light colored on the inner half, with the median area dusted with black. There are two strong raised scale tufts, and a more or less distinct notch opposite the cell. Brilliant metallic scaling is present along the outer margin below the notch, and a single row of four black dots is present along the outer margin. The foreleg is white with brown banding, while the other legs are predominantly white.

DISTRIBUTION: *Dicymolomia julianalis* is found throughout much of the eastern US and adjoining areas of southern Canada (Ontario; Quebec; New Brunswick; Nova Scotia). In the eastern US, the range extends from Maine to southern Florida, and westward to central Texas, central Oklahoma, central Kansas, Iowa, and Minnesota. In North Carolina, populations occur from the Coastal Plain westward to the lower elevations in the mountains.

FLIGHT COMMENT: The adults are active nearly year-round in Florida, and mostly from April through November in other areas outside of North Carolina. As of 2023, our records extend from early April through early November. Populations in the southern portions of the range presumably have more than one generation per year. (Landau et al., 1995) reported that there are three generations per year in Tennessee. The flight season in North Carolina is rather long and presumably reflects two or more overlapping generations per year.

HABITAT: In North Carolina, this species is associated with sites that support cattails such as farm ponds, ditches, and marshes.

FOOD: Broadleaf Cattail (*Typha latifolia*) appears to be the most widely used host, and is the only documented host in North Carolina as of 2023. Other species appear to serve as minor or secondary hosts (Monroe, 1972; Landau et al., 1996; Robinson et al., 2010). These include Musk Thistle (*Carduus nutans*), LeConte's Thistle (*Cirsium lecontei*), Canadian Milkvetch (*Astragalus canadensis*), a prickly-pear (*Opuntia* sp.), and a South American ornamental (*Amaranthus caudatus*). The larvae also have been found in dead bolls of cotton (*Gossypium* sp.) and have been observed feeding on clusters of bagworm eggs (*Thyridopteryx ephemeraeformis*). They also appears to function as endoparasitoids of larvae and pupae of bagworms (Powell et al., 1992).

OBSERVATION_METHODS: The adults are attracted to lights. The larvae can be found by breaking apart cattail heads during the winter and spring months and searching for the larvae. The larvae of *Limnaecia phragmitella* are also often present in cattail heads, but can be easily distinguished by their striped abdomen.

NATURAL HERITAGE PROGRAM RANKS: GNR S4S5

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: This species appears to be secure given that it is found statewide and its most common host (*Typha*) is widespread and common throughout the state.