




Bucculatrix inusitata No common name



| Status | Rank |
|--------|--------|
| NC | US |
| NC | Global |



FAMILY: Bucculatricidae SUBFAMILY: TRIBE:

TAXONOMIC_COMMENTS: *Bucculatrix* is a large genus of small leaf-mining moths, with around 300 species worldwide. A total of 103 Nearctic species have been described, and many others will likely be described in the future. Braun (1963) covered 99 species in her monograph, and four additional Nearctic species have been described since then.

FIELD GUIDE DESCRIPTIONS:

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS: Braun (1963)

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS: Braun's (1963) monograph on North American species of *Bucculatrix* remains the authoritative work on this group. The following is based on her study of specimens from throughout the range of this species. The head is white and the tuft may have a few brownish hairs. The antennal eye-cap is white, while the stalk is pale grayish ochreous and darker towards the tip. The thorax is white. The ground of the forewing is lustrous white and the markings ochreous. From just beyond the base to one-third the wing length, there is a pale longitudinal streak that is often faint or absent. Beginning at the basal third of the costa, there is an oblique streak that curves outward below the costa and narrows. It usually meets a second oblique costal streak behind it. This streak also narrows before extending as a narrow line to a small group of black-tipped raised scales on the termen near the tornus. At two-thirds, there is an irregular spot of variable size, with dark-tipped scales. These often encroach upon a triangular white area immediately before the apex and partially in the cilia. This triangular area extends across the wing to the group of dark-tipped raised scales near the tornus. Along its outer margin the smooth lustrous white scales of the general ground color form a narrow, almost iridescent bar lying alongside of the black-tipped scales which margin the termen. A second line of scales is present in the cilia, which are white before the apex, fuscous tinged opposite the apex, and white at the tornus. Their dark tips are at about the basal third of the cilia and curve inward near the apex toward the terminal line of black-tipped scales. From the middle of the dorsum there is a more or less distinct oblique streak that curves to the middle of the wing. The hindwings and cilia are grayish ochreous, while the legs are dull ochreous and shaded with fuscous. Braun (1963) notes that the best distinguishing wing characters of this species are the pure white triangular costal area before the apex, and the narrow lustrous white bar along the termen. The latter stands out sharply when light strikes it at an angle.

DISTRIBUTION: *Bucculatrix inusitata* is primarily found in the northeastern US and adjoining areas of Canada. Braun (1963) documented specimens from Quebec, Ontario, Maine, Massachusetts, New York, New Jersey, and Pennsylvania. MPG shows one record from Indiana. As of 2021, we have a single record from a high elevation site in Avery Co.

FLIGHT COMMENT: The adults fly during May, June and July in areas outside of North Carolina. Our one record as of 2021 is from 8 June.

HABITAT: The hosts are unknown and the habitat requirements are undocumented.

FOOD: There is one record of this species being raised from Common Juniper (*Juniperus communis*) based on a museum label, but Braun (1963) questioned the validity of this since this species belongs to a group of *Bucculatrix* that are either stem borers or gall formers on composites.

OBSERVATION_METHODS: The adults appear to rarely visit lights.

NATURAL HERITAGE PROGRAM RANKS: GNR SU

STATE PROTECTION:

COMMENTS: As of 2021 we currently have only one record for the state. Additional data are needed on the distribution and abundance of this species before we can assess its conservation status.