




*Crambus laqueatellus* Eastern Grass-veneer Moth



FAMILY: Crambidae SUBFAMILY: Crambinae TRIBE: Crambini  
 TAXONOMIC COMMENTS: One of 42 species in this genus that occur in North America north of Mexico (Scholtens and Solis, 2015), eleven of which have been recorded in North Carolina

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS: Fernald (1896); Forbes (1923)

TECHNICAL DESCRIPTION, IMMATURE STAGES: Fernald (1896)

ID COMMENTS: The ground color of the forewings is ochraceous with two longitudinal silver stripes separated by a strip of brown; a smaller spindle-shaped silver patch is also located between the outer silver streak and the costa. The area between the inner silver stripe and the inner margin is pale yellowish with longitudinal fuscous streaks and the lower part of the subterminal space is crossed by dark lines ((Fernald, 1896; Forbes, 1923)

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT:

FOOD:

OBSERVATION\_METHODS:

NATURAL HERITAGE PROGRAM RANKS:

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: