



Acrobasis cirroferella No common name



High counts of:
 1 - Carteret - 2019-08-26
 1 - Carteret - 2020-05-30
 1 - Carteret - 2021-09-06

● = Sighting or Collection in NC since 2004
 ○ = Not seen since 2004

Status	Rank
NC	US
NC	Global



FAMILY: Pyralidae SUBFAMILY: Phycitinae TRIBE: Phycitini
 TAXONOMIC_COMMENTS:

FIELD GUIDE DESCRIPTIONS:

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS: Neunzig (1996)

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS: "All *Acrobasis* with a 3 are very tricky for ID. In some cases it requires close look at male antennae, in others a look at the underside of the wings in males. In some cases females cannot be distinguished (e.g. *A. angusella* and *A. latifasciella*)" (Scholtens, 2017).

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: The habitat at the one site where this moth has been recorded in North Carolina supports extensive, dense thickets of Common Waxmyrtle.

FOOD: Larvae are reported to feed on Common Waxmyrtle (*Morella cerifera*) (Robinson et al., 2010). Tracy Feldman found a larva on *M. cerifera* in Scotland County believed to be this species (BugGuide record).

OBSERVATION_METHODS:

NATURAL HERITAGE PROGRAM RANKS: GNR SU

STATE PROTECTION:

COMMENTS: