



Homoeosoma asyloonnastes No common name




High counts of:

- 5 - Swain - 2004-07-19
- 2 - Swain - 2004-07-19
- 1 - Haywood - 2002-07-23

● = Sighting or Collection in NC since 2004
 ● = Not seen since 2004

Status	Rank
NC	US
NC	Global



FAMILY: Pyralidae SUBFAMILY: Phycitinae TRIBE: Phycitini
 TAXONOMIC_COMMENTS:

FIELD GUIDE DESCRIPTIONS:

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS: Neunzig (1996)

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS: This species closely resembles *Homoeosoma deceptorium*, but differs in being smaller (forewing length = 7.5-9.9 mm versus 10.0-12.5 mm for *H. deceptorium*). The black patch that follows the whitish antemedial line also tends to be more prominent on *H. deceptorium*. The males of *H. asyloonnastes* lack a thorn-like process on abdominal segment 8 that is present in *H. deceptorium* (Goodson and Neunzig 1993).

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT:

FOOD:

OBSERVATION_METHODS:

NATURAL HERITAGE PROGRAM RANKS:

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: