




Glena cribrataria Dotted Gray



FAMILY: Geometridae SUBFAMILY: Ennominae TRIBE: Boarmiini
TAXONOMIC_COMMENTS:

FIELD GUIDE DESCRIPTIONS: Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS: The ground color of the wings is pale whitish gray (Rindge, 1965). The lines consist of prominent black spots along the veins, with those on the postmedian and terminal most complete; the antemedian and median lines are represented mainly by dots on the costa, cubital and anal veins. A series of paired black dots runs down the dorsal surface of the abdomen.

DISTRIBUTION: Occurs statewide in North Carolina.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Our records come from both forests, heathlands, and other habitats with woody species.

FOOD: Larvae are broadly polyphagous on trees and shrubs. Wagner et al. (2001) specifically list maple (*Acer*), serviceberry (*Amelanchier*), birch (*Betula*), spruce (*Picea*), poplar (*Populus*), cherry (*Prunus*), oak (*Quercus*), willow (*Salix*), and blueberry (*Vaccinium*). The specific host plants used in North Carolina have not been recorded.

OBSERVATION_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 SNR [S4S5]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: