
Ectropis crepuscularia Small Engrailed Moth



FAMILY: Geometridae SUBFAMILY: Ennominae TRIBE: Boarmiini TAXONOMIC COMMENTS: Ectropis is a large, mainly paleotropical genus. Only one species occurs in our area.

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012) ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS: The ground color is pale gray. The lines are black, with both the antemedian and postmedian dentate. A strong double-tooth mark, including two points on the postmedian and two just following are usually diagnostic of this species (Forbes, 1948).

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: The vast majority of our records come from forests, almost all hardwoods bur with a few records from the summit of Mt. Mitchell, where spruce-fir forests dominate

FOOD: Larvae feed on a wide variety of woody plants, including some conifers in addition to hardwoods. Wagner et al. (2011) specifically list firs (<i>Abies</i> spp.), maples (<i>Acer</i> spp.), birches (<i>Betula</i> spp.), Sweetfern (<i>Comptonia peregrina</i>), ashes (<i>Fraxinus</i> spp.), Witch-hazel (<i>Hamamelis virginiana</i>), walnuts (<i>Juglans</i> spp.), larches (<i>Larix</i> spp.), apples (<i>Malus</i> spp.), spruces (<i>Picea</i> spp.), poplars (<i>Populus</i> spp.), cherries (<i>Prunus</i> spp.), Common Pear (<i>Pyrus communis</i>), oaks (<i>Quercus</i> spp.), currants (<i>Ribes</i> spp.), blackberries (<i>Rubus</i> spp.), willows (<i>Salix</i> spp.), hemlocks (<i>Tsuga</i> spp.), and Cranberry (<i>Vaccinium macrocarpon</i>). In North Carolina, larvae have been recorded on Box-elder (<i>Acer negundo</i>) and White Ash (<i>Fraxinus americana</i>).

OBSERVATION METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 SNR [S5]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: