




Hypagyrtis esther Esther Moth



FAMILY: Geometridae SUBFAMILY: Ennominae TRIBE: Bistonini

TAXONOMIC COMMENTS: This New World genus contains 7 species, 1 neotropical and 6 from North America, three of which occur in North Carolina (*piniata* may also be possibly present in the mountains -- Wagner et al., 2001). This genus shows a large range of variation, however, and its taxonomy still appears to be unsettled. Forbes (1948) stated that "the species or forms of this genus are completely confused and show no satisfactory differences in genitalia -- they may be merely races and strains, but breeding from known foods will be needed to make sure." Much of the variation is contained within *Hypagyrtis unipunctata* but there are forms believed to belong to that species that are highly similar to those of *esther*.

FIELD GUIDE DESCRIPTIONS: Covell (1984)

ONLINE PHOTOS: MPG, Bugguide

TECHNICAL DESCRIPTION, ADULTS: Forbes (1948)

TECHNICAL DESCRIPTION, IMMATURE STAGES: Wagner (2005; but possibly refers more directly to *H. piniata*)

ID COMMENTS: We follow the description given in Forbes (1948) that *Hypagyrtis esther* is distinguished by its even dark violet-brown ground color out to the subterminal line, followed by a rusty brown shade in the subterminal area. The antemedian and postmedian are both dark with the postmedian waved and most distinct; the postmedian is concave in its lower portion and meets the inner margin at a right angle. The subterminal pale spot typical of the genus is typically single and oval. Although not explicitly stated by Forbes, there is no indication of sexual dimorphism with respect to coloration and markings. Males are smaller and have proportionately shorter wings and are easily recognized by their pectinate antennae; the margins of the females are more scalloped. *Hypagyrtis brendae* is similarly dark but grayer, and has a similar rusty subterminal shading, but has a more conspicuous if diffuse median line. In *brendae*, the postmedian converges with the median towards the inner margin, with both lines following a straighter, more oblique course than the postmedian in *esther* (Heitzman, 1974). Covell (1984) states that there are melanic forms of *unipunctata*, which may resemble *esther*, but we have not seen any description of forms of *esther* that are buff, yellow-brown, or pale luteous colored as described by Forbes (1948) for the various forms now lumped as *unipunctata*, or the pepper-and-salt colored forms that he described for *H. piniata* and one form of *unipunctata*. Unfortunately, this genus does not appear to show any genitalic characters that can be used to distinguish the species (Forbes, 1948). Barcoding and other DNA analysis will probably be needed to resolve these issues. Until then, we recognize *esther* solely based on its dark, violet-brown median and basal areas and exclude buff or other more lightly colored or mottled forms.

DISTRIBUTION: Due to possible confusion with either an undescribed sibling of *esther* or with a form of *piniata*, the true distribution of *esther* is unclear in North Carolina. Forbes (1948) indicated that he only knew of specimens of *H. esther* from the Pine Barrens of New Jersey and the Coastal Plain of North Carolina. Wagner's (2005) account is also based on the assumption that *esther* is more likely to occur in the Coastal Plain and barrens communities and *piniata* more likely in the Mountains. Barcoding and additional host plant analysis is needed to clarify the situation.

FLIGHT COMMENT: While the species has been recorded throughout the growing season, there are probably three distinct broods in the Coastal Plain, two broods in the Piedmont and Low Mountains, and a single brood in the High Mountains. Second and third brood specimens are much smaller and somewhat different in maculation from the first brood.

HABITAT: In the Coastal Plain, we have records from habitats dominated by all three of the major pine species in that region: Longleaf Pine-dominated Savannas, Flatwoods, and Sandhills; Pond Pine Woodlands; and Loblolly-dominated Coastal Fringe Evergreen and Bottomland communities. In the eastern Piedmont, we have records from Xeric Hardpan Barrens where Short-leaf Pine and Virginia Pine are both common.

FOOD: Stenophagous, feeding on pines. We have reared it from larvae found on Loblolly Pine but it is not limited to the range of Loblolly. Possibly feeding on a wide range of pines but records are not available for most species.

OBSERVATION METHODS: Adults have short, non-functional mouthparts (Forbes, 1948); consequently, they do not come to bait or show up at flowers. They appear to come fairly well to blacklights with a maximum of 41 captured in a single trap.

NATURAL HERITAGE PROGRAM RANKS: G5 S5

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: Occupies a wide range of pine-containing habitats and probably feeds on a variety of pine species, including the widely planted Loblolly. Appears to be secure in the state, although more needs to be learned about its taxonomic status and true distribution in the state.