




Cepphis armataria Scallop Moth



FAMILY: Geometridae SUBFAMILY: Ennominae TRIBE: Anagolini
 TAXONOMIC_COMMENTS:

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Forests, woodlands, and wooded swamps (Wagner et al., 2001). Most of our records come from high elevation hardwoods, cove forests, and montane riparian forests.

FOOD: Larvae are polyphagous, feeding on many species of hardwood trees and shrubs. Wagner et al., 2001 specifically mention maple (<i>Acer</i>), birch (<i>Betula</i>), apple (<i>Malus</i>), and oak (<i>Quercus</i>). We do not have any feeding records in North Carolina.

OBSERVATION_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 SNR [S3S4]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: