




Idaea furciferata Notch-winged Wave Moth



FAMILY: Geometridae SUBFAMILY: Sterrhinae TRIBE: Sterrhini

TAXONOMIC_COMMENTS: One of thirty species in this genus that occur in North America north of Mexico (Pohl et al., 2016). Thirteen have been recorded in North Carolina.

FIELD GUIDE DESCRIPTIONS: Covell (1984)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Our records come mainly from fairly dry, open habitats. In the Coastal Plain, we have records from sandhill habitats; in the Piedmont, we have records both from mafic barrens, glades, Piedmont Longleaf glades, and dry monadnock forests; in the Mountains, our records come from lakeshore and residential areas.

FOOD: Host plants appear to be unknown

OBSERVATION_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G4 [S3S4]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: