




Idaea obfusaria Rippled Wave



FAMILY: Geometridae SUBFAMILY: Sterrhinae TRIBE: Sterrhini
 TAXONOMIC_COMMENTS: One of thirty species in this genus that occur in North America north of Mexico (Pohl et al., 2016). Thirteen have been recorded in North Carolina.

FIELD GUIDE DESCRIPTIONS: Covell (1984)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Our records come from both open, herb-dominated habitats, including savannas, flatwoods, sandridges, and bogs, and hardwood forests, including bottomlands, swamp forests, and dry ridges.

FOOD: Larvae are polyphagous, feeding on dead and live plant tissues of many trees, shrubs, and forbs (Wagner et al., 2001). We do not have any feeding records in North Carolina.

OBSERVATION_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G4G5 SNR [S4S5]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: