




# *Cyclophora packardi* Packard's Wave



*n=7*  
High Mt.



*n=64*  
Low Mt.




NC counties: 58

High counts of:


- 65 - Stokes - 1996-09-10
- 50 - Stokes - 1998-08-18
- 50 - Stokes - 1998-08-18

● = Sighting or Collection in NC since 2006  
● = Not seen since 2006

Status	Rank
NC	US
NC	Global



*n=206*  
*Pd*



*n=157*  
*CP*

FAMILY: Geometridae SUBFAMILY: Sterrhinae TRIBE: Cosymbiini  
TAXONOMIC\_COMMENTS:

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS: The ground color of the wings is buff, dusted with brown or red; the discal spot is white, surrounded by a heavy black border (Forbes, 1948).

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Our records come from a wide range of habitat types but mainly from wet to mesic forests.

FOOD: Larvae feed on oaks (Wagner et al., 2001). As of 2024, our only feeding record is for Turkey Oak (<i>Quercus laevis</i>).

OBSERVATION\_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G4 SNR [S5]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: