
Scopula limboundata Large Lace-border Moth



FAMILY: Geometridae SUBFAMILY: Sterrhinae TRIBE: Scopulini TAXONOMIC_COMMENTS: One of 26 species in this genus that occur in North America north of Mexico (Pohl et al., 2016), seven of which have been recorded in North Carolina

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012) ONLINE PHOTOS: TECHNICAL DESCRIPTION, ADULTS: Covell (1970) TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT:

FOOD: Larvae are polyphagous, feeding on many species of hardwood trees and shrubs, as well as forbs. Wagner et al. (2011) specifically list chokeberry (<i>Aronia</i>), Coastal Sweet-pepperbush (<i>Clethra alnifolia</i>), smartweed (<i>Persicaria</i>), cherry (<i>Prunus</i>), elm (<i>Ulmus</i>), and blueberry (<i>Vaccinium</i>).

OBSERVATION METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 SNR [S5]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: