



Hydriomena divisaria Black-dashed Hydriomena



FAMILY: Geometridae SUBFAMILY: Larentiinae TRIBE: Hydriomenini

TAXONOMIC COMMENTS: One of 56 species in this genus that occur in North America north of Mexico (McDunnough, 1954, covered 55 and Rindge added Hydriomena peratica in 1956). Most are boreal or are found in the West, with only five recorded in North Carolina.

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS: A medium-sized Wave. The ground color of the typical form (subspecies *divisaria* of McDunnough, 1954) is a dull, light gray

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Our records come primarily from high elevation forests but with a few from cove forests at lower elevations

FOOD: Reported hosts include fir (*Abies*), Eastern Hemlock (*Tsuga canadensis*), larch (*Larix*), spruce (*Picea*), and possibly pines (*Pinus*) (Maier et al., 2011).

OBSERVATION_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 S2S3

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: