




*Disclisioprocta stellata* Somber Carpet



FAMILY: Geometridae SUBFAMILY: Larentiinae TRIBE: Xanthorhoini  
TAXONOMIC\_COMMENTS:

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Found in North Carolina primarily in open habitats, from maritime dunes, inland sand ridges, old fields, and residential areas. A few also come from forested habitats.

FOOD: Forbes (1948) lists *Pisonia* (Devil's Claw) as a host plant, but that species is not found north of Florida. Covell (1984) adds Amaranths, one species of which -- Smooth Pigweed (*Amaranthus hybridus*) -- is found throughout North Carolina. Robinson et al. (2010) also list Marvel-of-Peru (*Mirabilis jalapa*) and American Pokeweed (*Phytolacca americana*), and there are multiple BugGuide records of larvae on pokeweed.

OBSERVATION\_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 S5

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: