




*Eupithecia broui* No common name



FAMILY: Geometridae SUBFAMILY: Larentiinae TRIBE: Eupitheciini  
 TAXONOMIC\_COMMENTS:

FIELD GUIDE DESCRIPTIONS:

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS: Rindge (1985)

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS: This species has broad forewings. The ground color is gray with numerous grayish brown scales (Rindge, 1985). A number of slender but clearly marked cross lines exist. The antemedian is angulate below the costa; an incomplete median line connects to the discal spot; and the curved postmedian is followed by pale shade line and a partial whitish subterminal. The basal segment of the abdomen tends to be slightly paler than the other segments.

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Flies in February, March, and April (Rindge, 1985)

HABITAT: Our records currently come from wet-mesic habitats but range from a wet pine savanna, a bay-lake shoreline, a mesic coastal bluff, and a montane alluvial forest.

FOOD: Larval stages and host plants are unknown (Rindge, 1985)

OBSERVATION\_METHODS:

NATURAL HERITAGE PROGRAM RANKS: GNR S2S3

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: