




Eupithecia swetti No common name



FAMILY: Geometridae SUBFAMILY: Larentiinae TRIBE: Eupitheciini
TAXONOMIC_COMMENTS:

FIELD GUIDE DESCRIPTIONS:

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS: McDunnough (1949)

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS: The forewings are elongate, more than twice as long as wide, but triangular in shape with an oblique outer margin (Forbes, 1948); the are pointed apically and have a checkered fringe (McDunnough, 1949). The ground color is light fawn, tending to gray (McDunnough, 1949). The markings are often obscure (McDunnough) or even, giving the appearance of striations (Forbes, 1948). The discal dot is slender. The second segment of the abdomen has a slight indication of a dark transverse band (McDunnough). *Eupithecia matheri* is similar in having narrow, elongated forewings and an early spring flight, but is grayer than *swetti* and has more obsolescent markings (Rindge, 1985). The hind wings are slightly paler than the forewings, and have a minute discal dot and may or may not have an extradiscal line or band (Rindge, 1985).

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: An early spring species

HABITAT:

FOOD: The early stages and food plant are apparently unknown (Rindge, 1985)

OBSERVATION_METHODS:

NATURAL HERITAGE PROGRAM RANKS: GNR S3S4

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: