




Callizzia amorata Gray Scoopwing



FAMILY: Uraniidae SUBFAMILY: Epipleminae TRIBE:
TAXONOMIC_COMMENTS:

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Our records come mainly from mesic hardwood forests but with one record from a rich levee forest. We do not have any records from dry sites with a mixture of pines, where Coral Honeysuckle can be common.

FOOD: Larvae feed on honeysuckles, including bush-honeysuckle (<i>Diervilla</i>) but perhaps not Japanese Honeysuckle (<i>Lonicera japonica</i>) (Wagner et al., 2001). Snowberry (<i>Symphoricarpos</i>) is used at least in the Pacific Northwest (Pacific Northwest Moths Website, accessed 2022-11-17). We do not have any feeding records in North Carolina.

OBSERVATION_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 SNR [S3S4]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: