




Callizzia amorata Gray Scoopwing Moth



FAMILY: Uraniidae SUBFAMILY: Epipleminae TRIBE:
TAXONOMIC_COMMENTS:

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Our records come mainly from mesic hardwood forests but with one record from a rich levee forest. We do not have any records from dry sites with a mixture of pines, where Coral Honeysuckle can be common.

FOOD: Larvae feed on Honeysuckles, including Bush Honeysuckle but perhaps not Japanese Honeysuckle (Wagner et al., 2001). Snowberry is used at least in the Pacific Northwest (Pacific Northwest Moths Website, accessed 2022-11-17).

OBSERVATION_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 S3S4

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: