




# *Tolype laricis* Larch Tolype



High counts of:

- 4 - Onslow - 1995-07-25
- 4 - Onslow - 1995-08-02
- 3 - Onslow - 1995-05-30

Status	Rank		
NC	US	NC	Global



*n=7*  
CP

Three periods to each month: 1-10 / 11-20 / 21-31

FAMILY: Lasiocampidae SUBFAMILY: Macromphaliinae TRIBE:  
TAXONOMIC COMMENTS:

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS: Franclemont (1973)

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS: Males can be distinguished from those of *T. notialis* by their smaller size and black color on the wings. Females are gray and cannot be distinguished from *T. notialis* based on maculation alone.

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT:

FOOD: Reported hosts include pine (*Pinus*), spruce (*Picea*), larch (*Larix*), and fir (*Abies*) (Franclemont, 1973; Beadle & Leckie, 2012).

OBSERVATION METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 SNR [SU]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: The identity of the dark forms of Tolype that are found in North Carolina -- including the Coastal Plain -- still need to be determined.