




Malacosoma disstria Forest Tent Caterpillar Moth



High counts of:

- 299 - Halifax - 2012-05-25
- 150 - Columbus - 2022-04-29
- 100 - Martin - 1996-05-22

Status	Rank
NC	US
NC	Global



FAMILY: Lasiocampidae SUBFAMILY: Lasiocampinae TRIBE: Lasiocampini
 TAXONOMIC_COMMENTS:

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Woodlands and Forests (Wagner, 2005). North Carolina records come from hardwood forests, ranging from Maritime Forests to Northern Hardwoods.

FOOD: Polyphagous, feeding on many hardwood trees and shrubs (Wagner, 2005). In swamp forests that are flooded during the winter, one of its natural control agents -- a Tachinid fly -- is greatly reduced and the populations of the moth can reach defoliating proportions. Water Tupelo is particularly heavily attacked, becoming stripped of leaves over 1000s of acres.

OBSERVATION_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 S5

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: