
Hyles lineata White-lined Sphinx



FAMILY: Sphingidae SUBFAMILY: Macroglossinae TRIBE: Macroglossini TAXONOMIC_COMMENTS: Our only truly cosmopolitan genus of some 29 species of which 3 occur in North America and 1 in North Carolina.

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012)

ONLINE PHOTOS: MPG, Bugguide, BAMONA

TECHNICAL DESCRIPTION, ADULTS: Forbes (1948); Hodges (1971); Tuttle (2007)

TECHNICAL DESCRIPTION, IMMATURE STAGES: Forbes (1948); Wagner (2005); Tuttle (2007)

ID COMMENTS: A striking, robust species. Fore-wings are dark brown with a tan diagonal streak and cross-banded with a series of fine white lines. The hindwings have bands of black along the costal and outer margins with a bright pink band in between. In our area, this species is likely to be confused only with the much larger <i>Eumorpha fasciata</i>, which has two tan streaks on the forewing and a pink outer edge to the hindwing. Sexes are similar.

DISTRIBUTION: Occurs statewide, from the Barrier Islands to the High Mountains.

FLIGHT COMMENT: Probably 2-3 broods over most of the state.

HABITAT: The majority of our records come from open habitats, especially dune habitats on Barrier Islands and lake and river shorelines and wet meadows inland.

FOOD: Stenophagous. Larvae feed on Portulaca and primroses; often large caterpillars on very small foodplants! Our few larval records all come from Evening Primrose.

OBSERVATION_METHODS: Actively visits flowers even during the day but does not come to bait. Appears to come relatively well to 15 watt UV lights and building lights. Larvae are often conspicuous and in the deserts of the west can form tremendous outbreaks.

NATURAL HERITAGE PROGRAM RANKS: G5 [S5]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: Statewide in distribution and occupying a wide range of open habitats, including disturbed areas, this species appears to be secure.