




*Pheosia rimosa* Black-rimmed Prominent



Status		Rank	
NC	US	NC	Global



FAMILY: Notodontidae SUBFAMILY: Notodontinae TRIBE: Notodontini  
 TAXONOMIC\_COMMENTS:

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Our one recent record for this species comes from a ridgetop site. Bigtooth Aspens occur at various sites within this county but are not known from the immediate vicinity of the collection site. Balsam Poplar (or more probably, Populus &times; jackii -- see Weakley, 2015) is common along the New River, the North Fork of which is located downslope from this site, but several miles away.

FOOD: Larvae feed mainly on Poplar but also on Willows (Wagner, 2005). Miller et al., (2018) state that the preferred hosts are Bigtooth Aspen, Trembling Aspen, and Balsam Poplar. Balsam Poplar has been recorded along the New River

OBSERVATION\_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 S1S2

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: