




## *Hyphantria cunea* Fall Webworm Moth



FAMILY: Erebiidae SUBFAMILY: Arctiinae TRIBE: Arctiini

TAXONOMIC COMMENTS: The only member of its genus in North America. Two species, *cunea* and *textor*, have been recognized in the past (e.g., Brimley, 1938; Forbes, 1960), but only *cunea* is considered valid now (Hodges, et al., 1983).

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012)

ONLINE PHOTOS: MPG, Bugguide, BAMONA

TECHNICAL DESCRIPTION, ADULTS: Forbes (1960)

TECHNICAL DESCRIPTION, IMMATURE STAGES: Forbes (1960); Wagner (2005)

ID COMMENTS: Adults are white with variable amounts of black spotting. They are typically smaller but similar to *Spilosoma* species, also possessing yellow on fore-coxae and femora like all but *S. latipennis*. Particularly similar to *S. congrua*, with both possessing all white abdomens and overlapping in size. The antennal shaft of *Hyphantria* is typically black, however (at least in the typical form, Forbes, 1960), whereas it is usually white in *Spilosoma* species (Forbes also describes form *textor* of *Hyphantria* as having white shafts). *Hyphantria* are also usually smaller, slimmer, and occasionally much darker than *Spilosoma*.

DISTRIBUTION: Occurs statewide (Brimley, 1938)

FLIGHT COMMENT: Has three broods in the Piedmont and Coastal Plain but the pattern is less clear in the Mountains

HABITAT: Wagner (2005) lists the habitats of this species as "yards and parks, fields, fencerows, woodlands, and forests." In North Carolina, it is found in most habitats that possess hardwood trees, including maritime and coastal fringe forests, sandhills, floodplains, mesic slopes, and dry ridges and summits.

FOOD: Polyphagous, but feeding solely on hardwood trees, unlike most other members of the Arctiinae. Recorded from over 400 species of hardwoods (Wagner, 2005). In North Carolina, we have observed it feeding on a wide range of hardwoods, including ash, birch, cherry, hickory, holly, maple, mulberry, persimmon, redbud, sourwood, sweetgum, sycamore, and walnut.

OBSERVATION\_METHODS: Populations are easily detected by the communal webs made by the larvae. Adults come well to blacklights, with up to 90 being collected in a single trap. None have been recorded at bait.

NATURAL HERITAGE PROGRAM RANKS: G5 [S5]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands

COMMENTS: This species is abundant, widespread, and makes use of a wide variety of habitats across the state, including wooded residential areas. It appears to be quite secure.