



Cycnia tenera Delicate Cycnia Moth



FAMILY: Erebiidae SUBFAMILY: Arctiinae TRIBE: Arctiini

TAXONOMIC COMMENTS: One of five species in this genus that occur in North America (Lafontaine and Schmidt, 2010), four of which have been recorded in North Carolina.

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012)

ONLINE PHOTOS: MPG, Bugguide, BAMONA

TECHNICAL DESCRIPTION, ADULTS: Forbes (1960)

TECHNICAL DESCRIPTION, IMMATURE STAGES: Forbes (1960); Wagner (2005)

ID COMMENTS: Fore-wings are white and somewhat translucent; head, thorax, and the costa of the forewing are deep yellow. Similar to *Cycnia inopinatus* but is larger, usually a purer white, and has the yellow costal stripe reaching well beyond the mid-point, often nearly to the apex of the forewing; legs are bicolored, unlike the solid lead gray in *inopinatus*. Most specimens should be easily distinguishable based on a good quality photograph.

DISTRIBUTION: Probably occurs statewide.

FLIGHT COMMENT: Wagner (2005) states that there are two to three generations per year; our data appear to show three peaks but with adults present essentially throughout the growing season.

HABITAT: Found in a wide variety of open habitats, including old field and road edges, where Dogbane is often abundant.

FOOD: Monophagous, feeding solely on Hemp Dogbane (*Apocynum cannabinum*).

OBSERVATION METHODS: Comes moderately well to blacklights and is often observed during the day, especially near patches of Dogbane. None of our records come from bait.

NATURAL HERITAGE PROGRAM RANKS: G5 [S5]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: Common, widespread, and uses a wide variety of habitats, including early successional and disturbed areas; appears to be quite secure.