




Zanclognatha theralis Flagged Zanclognatha



FAMILY: Erebidae SUBFAMILY: Herminiinae TRIBE:

TAXONOMIC COMMENTS: One of thirteen species in this genus that occur in North America north of Mexico (Lafontaine and Schmidt, 2010; 2013; Wagner and McCabe, 2011), all of which have been recorded in North Carolina

FIELD GUIDE DESCRIPTIONS:

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Our records come primarily from forested sites, but including both pine and hardwood dominated stands. Few, if any, come from open habitats, including savannas, maritime grasslands, and old fields.

FOOD: Larvae have been reared from Old Man's Beard (<i>Usnea</i>) (Sigal, cited by Wagner et al., 2011), but other members of this genus typically feed on dead leaves.

OBSERVATION_METHODS: Comes to blacklights.

NATURAL HERITAGE PROGRAM RANKS: G4 SNR [S4S5]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: