



Phalaenostola larentioides Black-banded Owlet Moth




High counts of:

100 - Stokes - 1998-08-18
100 - Stokes - 1996-09-10
50 - Stokes - 1998-08-18

● = Sighting or Collection in NC since 2006
● = Not seen since 2006

Status Rank
NC US NC Global



FAMILY: Erebiidae SUBFAMILY: Herminiinae TRIBE:
TAXONOMIC_COMMENTS:

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES: Wagner et al. (2011)

ID COMMENTS:

DISTRIBUTION: Occurs statewide in North Carolina.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Our records come from a wide range of woody and herbaceous, wet and dry, rich and nutrient-poor habitats.

FOOD: Larvae feed on dead and living grasses, as well as other detritus (Wagner et al., 2011).

OBSERVATION_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 SNR [S4S5]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: