




*Renia new species near discoloralis* No common name



High counts of:  
 5 - Orange - 2023-06-18  
 3 - Orange - 2023-05-23  
 3 - Orange - 2023-06-18

● = Sighting or Collection in NC since 2004  
 ● = Not seen since 2004

Status	Rank
NC	US
NC	Global



FAMILY: Erebiidae SUBFAMILY: Herminiinae TRIBE:  
 TAXONOMIC\_COMMENTS:

FIELD GUIDE DESCRIPTIONS:

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS: Forbes (1953); NatureServe Explorer (accessed 7-2020)

TECHNICAL DESCRIPTION, IMMATURE STAGES: Apparently undescribed

ID COMMENTS: This species is very similar to *Renia discoloralis* in its range of color and pattern phenotypes, but is noticeably smaller and flies in the spring and fall instead of mid-summer (Forbes, 1954).

DISTRIBUTION: Forbes (1954) believed that this was more of a southern species than true *discoloralis*. Schweitzer (in NatureServe Explorer) describes its range as extending as far north as New Jersey and Pennsylvania, but only names North Carolina as having definite occurrences in the South.

FLIGHT COMMENT: Adults are present as early as May in North Carolina. Records from September and October for members of this complex also probably represent this species

HABITAT: Our records come primarily from wet to mesic stands of hardwoods

FOOD: Larvae do not appear to have been observed but they probably feed on dead leaves and other detritus like other members of this genus

OBSERVATION\_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G4 S4

STATE PROTECTION:

COMMENTS: