




Hypena manalis Flowing-line Snout



FAMILY: Erebiidae SUBFAMILY: Hypeninae TRIBE:

TAXONOMIC COMMENTS: One of 29 species in this genus that occur in North America north of Mexico (Lafontaine and Schmidt, 2010). Fifteen species have been recorded in North Carolina.

FIELD GUIDE DESCRIPTIONS: Covell (1984; as Bomolocha manalis); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: The majority of our records come from bottomlands, swamps, and other wet habitats where False Nettle occurs. However, we also have several records from cove forests and stands of northern hardwoods, where other species of nettles are common.

FOOD: Larvae feed on False Nettle (*Boehmeria cylindrica*) (Wagner et al., 2011).

OBSERVATION_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 S4S5

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: