




## *Hypena baltimoralis* Baltimore Snout



FAMILY: Erebidae SUBFAMILY: Hypeninae TRIBE:

TAXONOMIC COMMENTS: One of 29 species in this genus that occur in North America north of Mexico (Lafontaine and Schmidt, 2010). Fifteen species have been recorded in North Carolina.

FIELD GUIDE DESCRIPTIONS: Covell (1984; as *Bomolocha baltimoralis*); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Occurs statewide

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT:

FOOD: Larvae feed on maples, including Red Maple (*Acer rubrum*) and Silver Maple (*A. saccharinum*) (Wagner et al., 2011). Our only feeding record in North Carolina is for Florida Maple (*A. floridanum*), but other maples are likely also used throughout the state.

OBSERVATION\_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 SNR [S5]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: