




## *Hypena eductalis* Red-footed Snout



FAMILY: Erebidae SUBFAMILY: Hypeninae TRIBE:

TAXONOMIC COMMENTS: One of 29 species in this genus that occur in North America north of Mexico (Lafontaine and Schmidt, 2010). Fifteen species have been recorded in North Carolina.

FIELD GUIDE DESCRIPTIONS: Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES: Wagner et al. (2011)

ID COMMENTS:

DISTRIBUTION: While we have records from the coast to the mountains, the species seems to be rare or absent from most of the Piedmont and Coastal Plain.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Our records come from a variety of river and pond shorelines, bogs, and other wetlands, particularly where alders are common

FOOD: Reported hosts include alder (*Alnus*) and River Birch (*Betula nigra*) (Wagner et al., 2011).

OBSERVATION\_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 SNR [S3S4]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: