
Anomis erosa Yellow Scallop Moth



FAMILY: Erebidae SUBFAMILY: Scoliopteryginae TRIBE: Anomini TAXONOMIC COMMENTS:

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS: Forbes (1954) TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS: This species is very similar in appearance to <i>Rusicada privata</i>. In contrast to <iR. privata</i>, <iA. erosa</i> has a well-developed tooth on both the antemedian and postmedian lines. In <iR. privata</i>, the antemedian line appears scalloped, without the obvious tooth of <iA. erosa</i> erosa</i> erosa</i> generally has a more yellowish ground color compared with <iR. privata</i> which can be heavily dusted with reddish scales. But there seems to be some overlap between redder specimens of <iA. erosa</i> and more yellow specimens of <iR. privata</i> , so coloration may not be a reliable means of distinguishing the species.

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Habitats are mainly open, including brushlands, croplands, marshes (Wagner et al., 2011).

FOOD: Larvae feed on many species in the Malvaceae, including cultivated species and native mallows (Wagner et al., 2011). Wagner (2005) specifically lists Hollyhock (<i>Alcea rosea</i>), Upland Cotton (<i>Gossypium hirsutum</i>) and <i>Hibiscus</i>). We do not have any feeding records in North Carolina.

OBSERVATION METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 SNR [S4S5]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: