



# *Mocis latipes* Small Mocis



High counts of:

23 - Brunswick - 1995-10-30

13 - Brunswick - 1995-10-23

12 - Brunswick - 1995-10-31

● = Sighting or Collection in NC since 2005  
● = Not seen since 2005

Status Rank  
NC US NC Global

FAMILY: Erebiidae SUBFAMILY: Erebinae TRIBE: Euclidiini  
TAXONOMIC\_COMMENTS:

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS: Strongly mottled and usually with a distinct median band and circular subreniform spot (Forbes, 1954)

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: The majority (well over 80%) of our records come from the Coastal Plain, including from maritime dunes, savannas, and sand ridges.

FOOD: Larvae feed on grasses, including crops such as corn, rice, sorghum, and sugarcane; turf grasses and pasture grasses (Wagner et al., 2011).

OBSERVATION\_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 SNR [S3S5]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: