




Catocala maestosa Sad Underwing



FAMILY: Erebiidae SUBFAMILY: Erebiinae TRIBE: Catocalini

TAXONOMIC COMMENTS: One of 103 species in this genus that occur in North America (Lafontaine and Schmidt, 2010, 2015), 67 of which have been recorded in North Carolina. Included by Barnes and McDunnough (1918) in their Group VI (also adopted by Forbes, 1954). This groups comprises 15 species, all of which feed on Hickories or Walnuts (Juglandaciae). In addition to maestosa, other members of this group that occur in North Carolina include retecta, dejecta, ulalume, insolabilis, myristica, vidua, lachrymosa, palaeogama, nebulosa, subnata, and neogama.

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: We have several records from urban areas, where planted Pecan may be the hosts. We also have a few records from brownwater river floodplains, where Bitternut or Water Hickory may be used.

FOOD: Larvae feed on Pecan and other Apocarya section hickories, as well as Walnut (Wagner et al., 2011)

OBSERVATION_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 S3S4

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: