
Catocala amestris Three-staff Underwing



FAMILY: Erebidae SUBFAMILY: Erebinae TRIBE: Catocalini

TAXONOMIC_COMMENTS: One of 103 species in this genus that occur in North America (Gall and Hawks, 2010; Kons and Borth, 2015a,b), 67 of which have been recorded in North Carolina.

FIELD GUIDE DESCRIPTIONS: Covell (1984)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION. ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES: Wagner et al. (2011)

ID COMMENTS: The ground color is brownish gray, with the basal and terminal areas usually solidly gray. The antemdian and postmedian lines are double and heavily black, forming a complex pattern of strong loops. The reniform also has a double black line around it but is often obscured by black shading (Sargent, 1976). The hindwing is a fairly deep yellow-orange and crossed by black bands. The inner band is sharply angled and the outer band can be either complete or broken.

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Our two records come from very different habitats, one from a xeric, coastal fringe sand ridge and the other from the foothills of the Blue Ridge.

FOOD: Larvae are stenophagous, feeding on <i>Amorpha</i> species. Wagner et al. (2011) list both False Indigo-bush (<i>Amorpha fruticosa</i>) and Leadplant (<i>A. canescens</i>). Locust (<i>Robinia</i>) has also been reported (Sargent, 1976), but Wagner discounts this claim. We do not have any feeding records in North Carolina.

OBSERVATION METHODS:

NATURAL HERITAGE PROGRAM RANKS: G4 SU [SH]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: