
Catocala lineella Little Lined Underwing



FAMILY: Erebidae SUBFAMILY: Erebinae TRIBE: Catocalini TAXONOMIC_COMMENTS: One of 103 species in this genus that occur in North America (Lafontaine and Schmidt, 2010, 2015), 67 of which have been recorded in North Carolina. Lineella was included by Barnes and McDunnough (1918) in their Group XX (also adopted by Forbes, 1954), which feed on Oaks. This group originally included amica and jair, with lineella now added as a separate species. At least one undescribed species may eventually be added. Catocala lineella has been treated as a form of amica, but was restored to full species status by Gall (1990), who also considered forms melanotica, aurantiaca, and novangliae of amica as a synonyms of lineella. Gall and Hawks (2010) treated form nerissa as an additional synonym of lineella.

FIELD GUIDE DESCRIPTIONS: Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES: Wagner et al. (2011)

ID COMMENTS: A medium-sized Underwing, with mottled, dark gray forewings and yellow-orange hindwings that lack a median black band. Grote (1872) described lineella as very similar to Catocala amica, but darker and "powdered with glaucous, bluish- or greenish-gray, and black scales". The curvifascia band that is present in some forms of amica -- at least as a well-defined dark arc rather than a diffuse shade -- is absent in lineella (Barnes and McDunnough, 1918; D.F. Schweitzer, pers. comm.). Catocala jair is usually more uniformly dark (in our populations) and has a straighter, less dentate postmedian line (Barnes and Mcdunnough, 1918).

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT:

FOOD: Larvae feed on oaks (<i>Quercus</i> spp.), particularly in the red and black oak groups (Wagner et al., 2011). Reported hosts include Scarlet Oak (<i>Q. coccinea</i>), Bear Oak (<i>Q. ilicifolia</i>), Chestnut Oak (<i>Q. montana</i>), Northern Red Oak (<i>Q. rubra</i>), Post Oak (<i>Q. stellata</i>), and Black Oak (<i>Q. velutina</i>) (Robinson et al., 2010). We do not have any feeding records in North Carolina.

OBSERVATION_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 SNR [S4S5]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: